

Phase-space approach to multi-population dynamics

Neil V. Budko · Fred J. Vermolen

the date of receipt and acceptance should be inserted later

Abstract Simultaneous deterministic dynamics of multiple populations described by a large set of ODE's is considered in the phase space of population sizes and ODE's parameters. The problem is formulated as a multidimensional phase-space conservation law and is solved explicitly for non-interacting multi-population models. Solutions for populations competing for a limited resource, populations with migration, as well as populations with phase-space interactions are obtained by simple iterative methods.

Keywords multi-population dynamics · phase-space distributions · migration · competition for limited resources

1 Introduction

Certain natural objects and phenomena, not immediately associated with population dynamics, can still be described as a large number of simultaneously developing populations. For example, forests consist of plants that in turn consist of many cells. Also cells themselves contain significant populations of mitochondria – small organelles capable of multiplying on their own. Thus, considering the mitochondria of a single cell to be the elementary population, one can view the cells of a plant as a population of populations, or a *multi-population*. Alternatively, viewing a single plant as an elementary population of cells, a field of plants is a population of populations, etc. (see, e.g., Merchant et al. 1960; Carrie et al. 2012).

Although, classical population dynamics is a very well developed subject (see, e.g., Shonwinkler and Herod 2009; Begon et al. 2006; Newman et al. 2014), its main concern has traditionally been the growth of a single or at most of a few populations simultaneously. The question of dynamics of dispersing

populations has led to the development of the metapopulation theory (e.g., Levin 1969; Eriksson et al. 2014) that treats populations with spatially disjoint habitats as a single metapopulation, and reduces the mathematical description to a single ODE. Alternatively, it is sometimes advantageous to consider the space-time dynamics of all members of all populations simultaneously, perhaps, subdividing them in a few species as well. This leads to spatially-resolved PDE models, such as the Fischer equation (Fischer 1930), that have produced many valuable insights over the years (see, e.g., Edelstein-Keshet 2005).

Studying multiple populations with large systems of ODE's or spatially-resolved PDE's does not always provide explicit answers to the pertaining questions. For example, in agriculture and ecology (e.g., Hautala and Hakojärvi 2011; Lin et al. 2013; Xu et al. 2014; Velazquez et al. 2014), where the growth of an individual plant or species is typically described by a logistic or a reaction-type ODE, the actual question of interest or the measured data is often the histogram of plant or population sizes. Predicting the time-evolution of such a histogram is not a trivial problem, since not only the initial conditions but also the various parameters of the ODE may differ from plant to plant. Another complication arises if populations interact in some way, e.g., there may be a competition for a limited resource or migration of individuals from one population to another. Populations may also interact on a more esoteric level, e.g., their growth rates may depend on the distribution over the phase space.

Histograms or their continuous analogue – phase-space distributions – are typically associated with random variables and stochastic processes and are less common in deterministic setting. Nonetheless, the temporal evolution of such deterministic distribution functions may be useful in the analysis of biological data, since it may help to distinguish deterministic causal effects from stochastic phenomena and random correlations. There is nothing new in the phase-space analysis of deterministic dynamics as such. In fact, it is often performed on the level of single trajectories (phase-space curves), e.g., the predator-prey system (see, e.g., Edelstein-Keshet 2005). The novelty of the present contribution is in the systematic extension of these ideas from single trajectories to distributions over the phase space.

To predict the time evolution of the multi-population distribution function we employ an n -dimensional phase-space conservation law, where the mathematical form of the phase-space current is determined by the dynamic equations (coupled or uncoupled ODE's) of individual populations. This approach has been inspired by the Lifshitz-Slyozov-Wagner method from materials science (Lifshitz and Slyozov 1961; Wagner 1961; Kampman and Wagner 1991; Myhr and Grong 2000; Collet and Goudon 2000; Laurencot 2002). Conceptually it may be viewed as a version of the Fokker-Planck equation tuned, in our case, to the specific needs of the multi-population dynamics. The present approach is also similar in spirit to the Liouville equation of Hamiltonian mechanics, except that the population dynamics is not necessarily Hamiltonian. The closest analogue in the field of population dynamics is, probably, the state-space method discussed in (Newman et al. 2014).

To illustrate the benefits and limitations of the proposed framework we investigate a range of population models relevant to plant biology and migration studies. We focus on problems that have either completely explicit solutions or can be solved with a simple iterative algorithm. In the subsequent sections we consider populations with unlimited (exponential) growth, populations whose growth is limited by the carrying capacity of the medium (logistic equation), populations competing for a single limited resource (e.g. oxygen, food, etc.), populations with migrating members, and populations whose rate of growth depends on the distribution function. Whenever possible we include both the population size and the parameters of ODE's in a single multidimensional distribution evolving in time.

2 Populations with unlimited growth

Let a very large number of populations be growing in accordance with the following equations:

$$\frac{dn_i}{dt} = \alpha_i n_i, \quad n_i(0) > 0, \quad i = 1, 2, \dots, P, \quad (1)$$

where $\alpha_i > 0$ are the growth constants. The variable n_i will be referred to as the size of the i -th population, but, in practice, it may denote many different things. For example, it can be the length, volume, or the biomass of a plant. It can also be a continuous approximation of the number of animals within a habitat, or the number of cells in a plant, or the number of mitochondria in a cell. The units of time are also completely arbitrary here.

Instead of solving all these ODE's we consider the time evolution of the distribution function $u(n, \alpha, t)$, whose integral over the size variable n and the growth constant variable α represents the total number of populations, which is conserved:

$$\int_0^\infty \int_0^\infty u(n, \alpha, t) dn d\alpha = P. \quad (2)$$

Integrating the distribution over α only,

$$\int_0^\infty u(n, \alpha, t) d\alpha = \rho(n, t), \quad (3)$$

one obtains a continuous approximation of the distribution of populations over their sizes at time t . The total size $N(t)$ of all populations at time t can be obtained as:

$$N(t) = \int_0^\infty n \rho(n, t) dn. \quad (4)$$

We assume that the sufficiently smooth continuous approximation of the initial distribution $u(n, \alpha, 0) = u_0(n, \alpha)$ is given. The latter is interpreted as a two-dimensional distribution of populations over their initial sizes n_0 and their growth constants α . Since the dynamics of all populations is purely deterministic, the functions u_0 and u are not probability density functions. However, upon suitable normalization, they can be interpreted as such for a randomly selected population.

The dynamics of populations causes the distribution function u to change in time. To account for these changes we introduce a phase-space current with the density $\mathbf{J}$. As a consequence of the conservation law (2) the distribution function u obeys the phase-space continuity equation:

$$\frac{\partial u}{\partial t} + \nabla \cdot \mathbf{J} = 0, \quad (5)$$

where the vector $\mathbf{J}$, in general, has as many components as there are phase-space variables, i.e. two in the present case:

$$\mathbf{J} = \langle uv_n, uv_\alpha \rangle. \quad (6)$$

The corresponding ‘velocities’ are deduced from the dynamic equation (1), i.e., in the present case: $v_n = n' = \alpha n$ and $v_\alpha = \alpha' = 0$. Since $\mathbf{J}$ has only one nonzero component, the continuity equation (5) reduces to the following Cauchy problem:

$$\frac{\partial u}{\partial t} + \alpha n \frac{\partial u}{\partial n} + \alpha u = 0, \quad u(n, \alpha, 0) = u_0(n, \alpha). \quad (7)$$

The solution of this problem, obtained with the method of characteristics, is

$$u(n, \alpha, t) = e^{-\alpha t} u_0(ne^{-\alpha t}, \alpha). \quad (8)$$

An example of the time evolution of $u(n, \alpha, t)$ is shown in Fig. 1, where the initial distribution $u_0(n, \alpha)$ is a Gaussian centered at $n_c = 15$ and $\alpha_c = 20$, which was set to zero for $\alpha, n < 0$ (see the top-left image). The size distributions of populations $\rho(n, t)$ (bottom row of Fig. 1) were obtained by numerical integration over α and demonstrate a progressively fattening tail due to the rapidly growing populations corresponding to the upper part of $u(n, \alpha, t)$, i.e., larger growth rate constants α . An interesting feature in the evolution of $u(n, \alpha, t)$ that becomes more pronounced at larger t ’s, is the gradual downwards shift of the maximum of $u(n, \alpha, t)$ with time. This is due to the fact that the populations with smaller α ’s tend to grow more evenly (synchronously) even if their initial n ’s are different, whereas, populations with larger α ’s and slightly different initial n ’s rapidly grow apart.




Fig. 1 Time evolution of $u(n, \alpha, t)$ according to (8). Top row – snapshots at $t = 0.0, 0.015$, and 0.03 ; Bottom row – corresponding numerically integrated $\rho(n, t)$ showing the distribution of populations over their sizes (the horizontal, n -axis, of these plots is shared with the images above).

3 Populations with logistic growth

Let the dynamics of populations be governed by the logistic equations

$$\frac{dn_i}{dt} = \gamma_i n_i (k_i - n_i), \quad n_i(0) > 0, \quad i = 1, 2, \dots, \quad (9)$$

where $\gamma_i \geq 0$ is a rate constant, and $k_i > 0$ is a constant determining the maximum sustainable population. The distribution function $u(n, \gamma, k, t)$ over the three-dimensional (n, γ, k) phase space satisfies the following Cauchy problem:

$$\frac{\partial u}{\partial t} + \gamma n(k - n) \frac{\partial u}{\partial n} + \gamma(k - 2n)u = 0, \quad u(n, \gamma, k, 0) = u_0(n, \gamma, k). \quad (10)$$

Here too the solution can be obtained with the method of characteristics. Since it is more involved than in the previous case, we shall provide the intermediate steps. The characteristic equations of (10) are

$$\frac{dt}{ds} = 1, \quad \frac{dn}{ds} = \gamma n(k - n), \quad \frac{du}{ds} = -\gamma(k - 2n)u. \quad (11)$$

This leads to the following relations:

$$C_0 = \frac{n}{|k - n|} e^{-\gamma k t}, \quad u = \frac{C_1}{n|k - n|}. \quad (12)$$




Fig. 2 Time evolution of $u(n, \gamma, k, t)$ according to (16) with fixed $k = 80$. Top row – snapshots at $t = 0.0, 0.001$, and 0.002 ; Bottom row – corresponding numerically integrated $\rho(n, t)$ showing the distribution of populations over their sizes.

We shall restrict ourselves to the segment $0 \leq n < k$, so that $|k - n| = k - n$. The general form of the solution to (10) in this case is

$$u = \frac{f(p)}{n(k-n)}, \quad p = \frac{n}{k-n} e^{-\gamma kt}. \quad (13)$$

The function $f(p)$ should be chosen in such a way that the initial condition is satisfied. To this end we introduce two smooth functions of p that at $t = 0$ behave as follows:

$$\left. \frac{kp}{p+1} \right|_{t=0} = n, \quad \left. \frac{k^2 p}{(p+1)^2} \right|_{t=0} = n(k-n). \quad (14)$$

The initial condition from (10) is satisfied if these functions are used as

$$\frac{k^2 p}{n(k-n)(p+1)^2} u_0 \left(\frac{kp}{p+1}, \gamma, k \right) \Big|_{t=0} = u_0(n, \gamma, k). \quad (15)$$

Thus, the solution of the Cauchy problem (10) on the strip $0 \leq n < k$ is

$$u(n, \gamma, k, t) = \frac{k^2 e^{-\gamma kt}}{(n e^{-\gamma kt} + k - n)^2} u_0 \left(\frac{k n e^{-\gamma kt}}{n e^{-\gamma kt} + k - n}, \gamma, k \right), \quad (16)$$

and for $n = k$, we obviously have $u(k, \gamma, k, t) = u_0(k, \gamma, k)$.

An example of the time evolution of $u(n, \gamma, k, t)$ for a fixed $k = 80$ is shown in Fig. 2, where the initial distribution $u_0(n, \gamma, 80)$ (top left) is the same Gaussian as in the example of Fig. 1. As t grows the corresponding

size distributions $\rho(n, t)$ (bottom row) tend to a distribution localized at the right boundary $n = k$. This time the maximum of the distribution $u(n, \gamma, k, t)$ shifts upwards (eventually), meaning that for larger t 's the populations with larger growth rates γ will dominate, as they will rapidly accumulate around the maximum attainable size.

4 Populations competing for a resource

Consider populations competing for a limited resource c that controls their growth rate. This control substance is absorbed equally well by all members of all populations. In response, the populations grow. However, we assume that the growth of a population slows down and eventually stops if the supply of c drops to some minimal value $c_{\min}$, which is here taken to be zero for simplicity. The following coupled system of equations describes this situation:

$$\begin{aligned} \frac{dc}{dt} &= -\gamma \sum_i n_i, \quad c(0) = c_0, \gamma > 0, \\ \frac{dn_i}{dt} &= \beta_i n_i c, \quad n_i(0), \beta_i > 0, \quad i = 1, 2, \dots \end{aligned} \quad (17)$$

Integrating the first equation and substituting the result in the second equation we arrive at:

$$\frac{dn_i}{dt} = \beta_i n_i \left(c_0 - \gamma \int_0^t \sum_j n_j(t') dt' \right), \quad i = 1, 2, \dots \quad (18)$$

The sum inside these equations is the total size $N(t)$ of all populations at time t , which in terms of the distribution function $u(n, \beta, t)$ can be written as

$$\sum_i n_i(t) = N(t) = \int_0^\infty \int_0^\infty n u(n, \beta, t) dn d\beta. \quad (19)$$

Thus, the phase-space dynamics of the distribution function will be governed by a weakly nonlinear equation:

$$\frac{\partial u}{\partial t} + \beta c n \frac{\partial u}{\partial n} + \beta c u = 0, \quad u(n, \beta, 0) = u_0(n, \beta), \quad (20)$$

where

$$c(t) = c_0 - \gamma \int_0^t N(t') dt'. \quad (21)$$

The questions of existence and uniqueness of the solutions for this and similar equations appearing below are treated in (Collet and Goundon 2000; Laurençot 2002).

As expected, the equation (20) confirms that the distribution u stops changing in time (becomes stationary) when the supply c is exhausted. Although, this behaviour is asymptotic for $t \rightarrow \infty$, one may expect that the changes in u are negligible for all $t > T$ for some sufficiently large T .

Let us focus on finding the solution u of the linear Cauchy problem (20) for a given function $c(t)$. Exploiting the direct analogy with (8) the solution of (20) is found to be

$$u(n, \beta, t) = e^{-\beta \xi(t)} u_0(n e^{-\beta \xi(t)}, \beta), \quad (22)$$

where

$$\xi(t) = \int_0^t c(\tau) d\tau. \quad (23)$$

Equation (22) is an implicit solution of the original problem, since $\xi(t)$ depends on u via (19), (21), and (23).

Let $u_k(n, \beta) = u(n, \beta, t_k)$, $N_k = N(t_k)$, $c_k = c(t_k)$, $\xi_k = \xi(t_k)$, and $t_{k+1} = t_k + \Delta t$. For sufficiently small Δt we can approximate the solution with the following time-stepping scheme:

Algorithm 4

- Given: $\xi_0 = 0$, $c_0 > 0$, $u_0(n, \beta)$, and a sufficiently large Ω .
- For $k = 0, 1, 2, \dots$, while $c_k \geq 0$, do:

$$\begin{aligned} \xi_{k+1} &= \xi_k + c_k \Delta t, \\ u_{k+1}(n, \beta) &= e^{-\beta \xi_{k+1}} u_0(n e^{-\beta \xi_{k+1}}, \beta), \\ N_{k+1} &= \iint_{\Omega} n u_{k+1}(n, \beta) dn d\beta, \\ c_{k+1} &= c_k - \gamma N_{k+1} \Delta t \end{aligned}$$

An example of the time evolution of $u(n, \beta, t)$ and the corresponding consumption of the resource $c(t)$ computed with the Algorithm 4 is shown in Fig. 3. Due to the mathematical equivalence of the two solutions, the states (snapshots) of $u(n, \beta, t)$ given by (22) are exactly equal to the states $u(n, \alpha, t)$ of the unmitigated exponential growth given by (8), provided $\alpha = \beta$. However, the evolution of $u(n, \beta, t)$ proceeds at a different rate and freezes as $c(t) \rightarrow 0$. The accuracy of the Algorithm 4 can be eventually improved using the predictor-corrector technique.

5 Populations with randomized migration

Let a single metapopulation be nonuniformly distributed over some spatial domain at time $t = 0$. The whole habitat may be divided either naturally or virtually into a large number of elementary cells (sub-habitats) and a question




Fig. 3 Initial distribution $u_0(n, \beta)$ (top-left), the final quasi-stationary distribution $u_0(n, \beta, t_{\max})$ (top-right), and the corresponding time-evolution of the resource $c(t)$ (bottom).

can be posed about the distribution of cell populations over their size at time t . The growth of these elementary cell populations could be described by one of the models from the preceding sections with growth parameters varying from cell to cell. If all individual members stay within their original cells or the whole metapopulation uniformly translates in space the phase-space methods developed earlier and the resulting distribution functions, obviously, apply without change.

Allowing for the migration of individuals between the cells usually requires a PDE-based space-time dynamic law (e.g. Fischer equation). Such a law cannot be directly incorporated into the phase-space current, since the phase-space approach completely neglects the spatial ordering of the cells. Nevertheless, certain types of migration mechanisms can be described on the level of ODE's and the corresponding phase-space continuity laws.

Consider a stationary metapopulation, where the total number of individuals and the total number of sub-habitats (that play the role of populations in this case) do not change. Suppose that all individuals from time to time (with frequency β) decide to emigrate from its native population so that each population is subject to the emigration rate βn , proportional to its size. Depending on the choice strategy of migrants as far as their new population is concerned one arrives at different ODE's.

For example, let each emigrant choose a new population (habitat) completely at random. If viewed spatially, this is, obviously, a variant of super-diffusion with a random step size. Then, on a sufficiently large time scale, there will be the steady immigration rate γ in all populations (since the total

emigration rate is constant). The emigration and immigration coefficients β and γ must be balanced to guarantee the steady state of the metapopulation:

$$\frac{d}{dt} \sum_{i=1}^P n_i = \sum_{i=1}^P (\gamma - \beta n_i) = 0. \quad (24)$$

Let $\sum_i n_i = N$, then $\gamma = \beta \bar{n}$, where $\bar{n} = N/P$, and P is the number of habitats. The dynamic law becomes

$$\frac{dn_i}{dt} = \beta (\bar{n} - n_i), \quad i = 1, 2, \dots, P. \quad (25)$$

Notice that $\bar{n} = N/P$ is the total number of individuals divided by the number of habitats, which would be exactly the size of populations in all habitats if all individuals were evenly spread throughout. From (25) it follows that a population stops changing once it reaches this equilibrium size $\bar{n}$.

The distribution function $u(n, t)$ is then found by solving the following Cauchy problem:

$$\frac{\partial u}{\partial t} + \beta (\bar{n} - n) \frac{\partial u}{\partial n} - \beta u = 0, \quad u(n, 0) = u_0(n), \quad (26)$$

and is given by

$$u(n, t) = \begin{cases} e^{\beta t} u_0(\bar{n} - (\bar{n} - n)e^{\beta t}), & n < \bar{n}, \\ e^{\beta t} u_0(\bar{n}), & n = \bar{n}, \\ e^{\beta t} u_0(\bar{n} + (n - \bar{n})e^{\beta t}), & n > \bar{n}. \end{cases} \quad (27)$$

The middle branch of this solution shows that the number (density) of populations with $n = \bar{n}$ growth exponentially with time. As can be seen in Fig. 4 the upper and lower branches of the solution (27) represent the initial distribution that shrinks, respectively, from the left and from the right, towards $\bar{n}$. This behaviour is indicative of a mollifier of the Dirac delta function centred at $\bar{n}$, which, indeed, is the limit of $u(n, t)$ as $t \rightarrow \infty$, since all populations will eventually have the same size $\bar{n}$. An interesting feature of (27) is the preservation of the main features of the initial $u_0(n)$ over time, albeit in a scaled (shrank) form, e.g., the double peaked shape shown in Fig. 4.

6 Biased migration

Consider populations with potentially unlimited growth and constant emigration frequency. Let the choice of the new population by the migrants be biased towards populations with higher growth rates. The ODE describing this situation is:

$$\frac{dn_i}{dt} = \alpha_i n_i - \beta n_i + \gamma \alpha_i, \quad i = 1, \dots, P, \quad (28)$$




Fig. 4 Time evolution of the distribution function $u(n, t)$ for populations with (uniform) migration given by the equation (27) with $\beta = 1.0$ and $\bar{n} = 5.7$.

where α_i is the growth coefficient, β is the emigration frequency, and γ is the immigration coefficient. Since migration does not change the total number of individuals (only the growth does), the following balance equation should be satisfied at all times:

$$\beta \sum_{i=1}^P n_i = \gamma \sum_{i=1}^P \alpha_i, \quad (29)$$

reducing equation (28) to

$$\frac{dn_i}{dt} = (\alpha_i - \beta)n_i + \alpha_i \beta \frac{\sum n_j}{\sum \alpha_j}, \quad i, j = 1, \dots, P. \quad (30)$$

The corresponding nonzero component of the phase-space current is

$$J_n = (\alpha - \beta)nu + \alpha\beta Ru, \quad (31)$$

where

$$R = \frac{N}{A}, \quad (32)$$

$$N = \iint nu(n, \alpha, t) d\alpha dn, \quad A = \iint \alpha u(n, \alpha, t) d\alpha dn,$$

and, since the α -component of the phase-space current is zero, A will stay constant in time. Thus, we arrive at the following weakly nonlinear continuity equation for the distribution $u(n, \alpha, t)$:

$$\frac{\partial u}{\partial t} + (\alpha\beta R + (\alpha - \beta)n) \frac{\partial u}{\partial n} + (\alpha - \beta)u = 0, \quad u(n, \alpha, 0) = u_0(n, \alpha). \quad (33)$$




Fig. 5 Time evolution of the distribution function $u(n, \alpha, t)$ for populations with biased migration, see eq. (34). Top row – snapshots at $t = 0.0, 0.018$, and 0.034 ; Bottom row – corresponding numerically integrated $\rho(n, t)$ showing the distribution of populations over their sizes.

Assuming that $R(t)$ is a given function of time we use the method of characteristics to obtain the implicit solution of this problem as

$$u(n, \alpha, t) = e^{-(\alpha-\beta)t} u_0 \left(n e^{-(\alpha-\beta)t} - \alpha \beta \xi(\alpha, t), \alpha \right),$$

$$\xi(\alpha, t) = \int_0^t R(t') e^{-(\alpha-\beta)t'} dt'. \quad (34)$$

However, $R(t)$ and, hence, $\xi(\alpha, t)$ are functions of u as well, see eq. (32). Thus, we employ an iterative algorithm similar to the Algorithm 4.

Algorithm 6

- Given: $\xi_0(\alpha) = 0$, $u_0(n, \beta)$, N_0 , R_0 , and a sufficiently large Ω .
- For $k = 0, 1, 2, \dots$, do:

$$\xi_{k+1}(\alpha) = \xi_k(\alpha) + R_k e^{-(\alpha-\beta)t_k} \Delta t,$$

$$u_{k+1}(n, \alpha) = e^{-(\alpha-\beta)t_{k+1}} u_0 \left(n e^{-(\alpha-\beta)t_{k+1}} - \alpha \beta \xi_{k+1}(\alpha), \alpha \right),$$

$$R_{k+1} = \frac{1}{A} \iint_{\Omega} n u_{k+1}(n, \alpha) dn d\alpha,$$

A quick look at the implicit solution (34) reveals similarity with the case of exponential growth (7). Indeed, as expected, the two solutions are identical for

$\beta = 0$, remain similar at low values of the emigration frequency, and become very different for larger values. One of the predictions of the metapopulation theory (Levin 1969; Eriksson et al. 2014) is that migration may prevent extinction. In the present case, as follows from (30), none of the populations can disappear as long as there is at least one nonzero population somewhere. Yet, it does not mean that all populations are always growing in size. In the early stages some populations may decrease. From (30) we deduce that the condition on the growth of a population is

$$\frac{\alpha_i}{\sum_j \alpha_j} > \left(1 - \frac{\alpha_i}{\beta}\right) \frac{n_i}{\sum_j n_j}, \quad \beta > 0. \quad (35)$$

Obviously, this condition is satisfied for all populations with $\alpha_i > \beta$. However, for a population with $\alpha_i < \beta$ it is easy to imagine a sufficiently large initial state $n_i(0)$ such that this condition is violated and the population size will decrease. Nevertheless, when such a decreasing n_i becomes sufficiently small, the condition will become satisfied again and the population will resume its growth.

An example of the evolution of distribution function for populations with biased migration computed with the Algorithm 6 is shown in Fig. 5. The initial distribution (the leftmost image and plot) is a Gaussian centred at $n_c = 20$ and $\alpha_c = 30$. The emigration frequency is set at $\beta = 50$ in this simulation, which is higher than most of the growth coefficients α in the initial distribution. The first striking conclusion is that despite the immigration bias we do not observe an explosive growth of some particular (chosen) populations at the expense of others. Apparently, any such growth is efficiently mitigated by a higher emigration rate.

The early-time transient decrease of some populations mentioned above (those with large initial n 's) causes the distribution function to contract slightly in the horizontal n -dimension, since the right side of the distribution moves to the left, while the left side moves to the right. This happens in the very early stages of the evolution (not shown). The same effect prevents the distribution from widening in the horizontal direction in the course of the evolution. In fact, as the Figure 5 shows, the distribution only keeps contracting.

Thus, the main consequence of the large emigration frequency β for the late-time evolution appears to be the uniformity of size for populations with equal α 's and the uniformity of α 's for populations of the same size. Also, for large t 's the populations show a very strong correlation (almost direct proportionality) between the population size and its growth coefficient α (see the top-right image of Fig. 5).

7 Growth rates depending on the distribution function

The present approach is indispensable for dynamic models where the rate of growth of populations is a functional $R(u, n)$ of both the size n of the local population and the distribution u of populations over the phase space. The

phase-space current vector will have as many nonzero components as there are time varying parameters in R . If all parameters, except n_i , are time invariant, then the problem is described by the following equations:

$$\frac{dn_i}{dt} = R(u, n_i), \quad i = 1, 2, \dots, P, \quad (36)$$

$$\frac{\partial u}{\partial t} = -\frac{\partial}{\partial n} (uR(u, n)). \quad (37)$$

Obviously, it is now impossible to solve the dynamic equations (36) without first solving the phase-space problem (37).

Consider populations with unlimited growth discussed earlier. Suppose that one is able to influence the rate of growth of each population (e.g., by judiciously watering and fertilizing each plant) proportionally to its current ‘weight’ in the phase space. Such a problem is described by the following equations:

$$\frac{dn_i}{dt} = \alpha_i n_i \int_{\alpha_i - \Delta\alpha}^{\alpha_i + \Delta\alpha} \int_{n_i - \Delta n}^{n_i + \Delta n} u(n, \alpha, t) dn d\alpha \approx \alpha_i n_i u(n_i, \alpha_i, t) \delta, \quad i = 1, 2, \dots, P, \quad (38)$$

$$\frac{\partial u}{\partial t} + 2\alpha\delta nu \frac{\partial u}{\partial n} + \alpha\delta u^2 = 0, \quad u(n, \alpha, 0) = u_0(n, \alpha), \quad (39)$$

where $\delta = 4\Delta\alpha \Delta n$, i.e., we use the mid-point approximation for the integral in (38). The resulting approximate phase-space problem (39) is a variant of the Burgers equation (see e.g. Smoller 1994) whose characteristic equations are:

$$\frac{dt}{ds} = 1, \quad \frac{dn}{ds} = 2\alpha\delta nu, \quad \frac{du}{ds} = -\alpha\delta u^2, \quad (40)$$

or

$$dn = 2\alpha\delta nu dt, \quad -\alpha\delta dt = \frac{du}{u^2}, \quad \frac{dn}{2n} = -\frac{du}{u}. \quad (41)$$

The solutions of the last two are:

$$\alpha\delta t - \frac{1}{u} = C_2, \quad u\sqrt{n} = C_3, \quad (42)$$

which allows rewriting and solving the first equation in (41) as

$$\frac{dn}{\sqrt{n}} = 2\alpha\delta C_3 dt, \quad 2\sqrt{n}(1 - \alpha\delta tu) = C_1. \quad (43)$$

Combining this result with the first equation in (42) we get

$$u = \frac{1}{\alpha\delta t - f(p)}, \quad p = 2\sqrt{n}(1 - \alpha\delta tu). \quad (44)$$

The initial condition requires that

$$f(p)|_{t=0} = -\frac{1}{u_0(n, \alpha)}. \quad (45)$$

Hence, we choose

$$f(p) = -\frac{1}{u_0(p^2/4, \alpha)} = -\frac{1}{u_0(n(\alpha\delta t u - 1)^2, \alpha)}, \quad (46)$$

and arrive at the following equation that implicitly defines the distribution $u(n, \alpha, t)$:

$$u = \frac{u_0(n(\alpha\delta t u - 1)^2, \alpha)}{\alpha\delta t u_0(n(\alpha\delta t u - 1)^2, \alpha) + 1}, \quad (47)$$

or

$$u = (1 - \alpha\delta t u) u_0(n(1 - \alpha\delta t u)^2, \alpha). \quad (48)$$

These representations are only valid in the smooth regime, i.e., prior to the development of shock.

The fixed-point iterative algorithm that computes the distribution at a given time t may be formulated as follows (one can use any suitable norm for the computation of the current mismatch ϵ_k):

Algorithm 7

- Given: $u_0(n, \alpha)$, t , and ϵ .
- For $k = 0, 1, 2, \dots$, while $\epsilon_k > \epsilon$, do:

$$u_{k+1}(n, \alpha) = \frac{u_0(n(\alpha\delta t u_k(n, \alpha) - 1)^2, \alpha)}{\alpha\delta t u_0(n(\alpha\delta t u_k(n, \alpha) - 1)^2, \alpha) + 1},$$

$$\epsilon_k = \|u_{k+1}(n, \alpha) - u_k(n, \alpha)\|.$$

Figure 6 illustrates the application of this algorithm (with $\epsilon = 10^{-4}$ and $\delta = 0.1$). The initial distribution is the same as in the example of the previous section. As expected from the non-viscous Burgers equation the solution develops a shock (notice the almost vertical front in the plot at the bottom-right of Fig. 6). Even though we have used the distribution computed at the previous time instant as the initial guess for the next time instant, the convergence of the Algorithm 7 slows down as time grows. Around $t = 0.1$, as one approaches the shock in this example, the convergence of the Algorithm 7 breaks down completely: initially the error ϵ_k decreases, however, it never reaches the tolerance of $\epsilon = 10^{-4}$ and even begins to increase as iterations continue.

Obviously the development of shock and the related breakdown of the Algorithm 7 does not mean that the populations stop growing. Rather, it is a result of the midpoint approximation of the integral in the equation (38). Keeping that in mind, we see that at least in the early stages of evolution there will emerge a sharp upper bound on the sizes of populations and many populations with the average growth factor will have that size.




Fig. 6 Time evolution of the distribution function $u(n, \alpha, t)$ for populations with growth rates depending on u , see eq. (47). Top row – snapshots at $t = 0.0, 0.054$, and 0.09 ; Bottom row – corresponding numerically integrated $\rho(n, t)$ showing the distribution of populations over their sizes.

8 Conclusions and possible extensions of the model

The phase-space approach to multi-population dynamics presented above allows computing the time evolution of the distribution function that approximates the multi-dimensional histogram of populations. Although we have considered some trivial stochastic processes in the sections about migration, the main value of this approach is in the fast and easy analysis of causal deterministic trends in biological data. For example, in the case of precision agriculture (Hautala and Hakojärvi 2011), this method could be employed to predict and, perhaps, control the distribution of plant sizes at the time of harvest. To this end, instead of introducing stochastic variations due to the environment, models considered above can be easily extended to include deterministic (measured) time-varying growth coefficients and environmental factors. Another straightforward extension could be the inclusion of the source term in the conservation law (5) reflecting the emergence of new populations.

The case of populations competing for limited resources could serve as a mathematical model for recovering the distribution of populations from the resource consumption curves. For example, in the analysis of seed quality it is common to monitor the oxygen consumption during the germination process (Bewley et al. 2013, van Duin and Koenig 2001, van Asbrouck and Taridno 2009), with the main consumers of oxygen being the growing populations of mitochondria present in every seed. Our method provides the neces-

sary link between the distribution of the seed parameters, including the initial number of active mitochondria, and the measured oxygen uptake curves for a batch of seeds germinating in a closed container.

It is easy to imagine that the phase-space dynamics would be even more exciting if the phase-space current had several nonzero components, e.g., with the time varying growth/environmental factors or with the classical predator-prey problem. The corresponding continuity equations, however, would have to be solved numerically (Leveque 2002).

Although we were able to analyze two special migration problems, the general case of linear coupling between ODE's so far appears to be intractable with our methodology. This has mainly to do with the nonlinearity of the mapping between a given vector and its histogram or distribution. Indeed, the histogram of the sum of two vectors is not the sum of the histograms of these vectors. Instead, we should, probably, focus on the phase-space coupling considered in the last section.

Acknowledgements and ethical statement

The authors are grateful to Prof. A. van Duijn (Leiden University and Fytagoras BV) and Dr. S. Hille (Leiden University) for introduction into the field of plant physiology and the problem of seed germination, and to Dr. H. van Doorn (HZPC Holland BV) for introduction to the problems of precision agriculture. The authors declare no conflicts of interest.

References

1. Begon M, Townsend CR, and Harper JL (2006) Ecology: From Individuals to Ecosystems. Blackwell Publishing.
2. Bewley JD, Bradford KJ, Hilhorst HWM, and Nonogaki H (2013) Seeds: Physiology of development, germination and dormancy. 3rd Edition, Springer, New York, Heidelberg, Dordrecht, London.
3. Carrie C, Murcha MW, Giraud E, Ng S, Zhang MF, Narsai R, and Whelan J (2012) How do plants make mitochondria? *Planta: An International Journal of Plant Biology*, 10.1007/s00425-012-1762-3.
4. Collet JF and Goudon T (2000) On solutions of the Lifshitz-Slyozov model. *Nonlinearity* 13:1239–1262.
5. Edelstein-Keshet L (2005) *Mathematical Models in Biology*. SIAM.
6. Eriksson A, Elias-Wolff F, Mehlig B, and Manica A (2014) The emergence of the rescue effect from explicit within- and between-patch dynamics in a metapopulation. *Proc. R. Soc. B* 281:20133127.
7. Fisher RA (1930) *The genetical theory of natural selection*. Oxford University Press.
8. Hautala M and Hakojärvi M (2011) An analytical C3-crop growth model for precision farming. *Precision Agric.* 12:266–279.
9. Kampman R and Wagner R (1991) *Materials Science and Technology - A Comprehensive Treatment*. Vol. 5, VCH, Weinheim, Germany.
10. Laurençot P (2002) The Lifshitz-Slyozov-Wagner equation with conserved total volume. *SIAM J. Math. Anal.* 34(2):257–272.
11. Leveque RJ (2002) *Finite volume methods for hyperbolic problems*. Cambridge University Press, Cambridge.

12. Levin R (1969) Some demographic and genetic consequences of environmental heterogeneity for biological control. *Bull. Entomol. Soc. Am.* 15:237–240.
13. Lifshitz IM and Slyozov VV (1961) The kinetics of precipitation from supersaturated solid solutions. *J. Phys. Chem. Solids*, 19:35–50.
14. Lin Y, Berger U, Grimm V, Huth F, and Weiner J (2013) Plant Interactions Alter the Predictions of Metabolic Scaling Theory. *PLoS ONE* 8(2):e57612.
15. Merchant DJ, Kuchler RB, and Munyo W (1960) Population dynamics in suspension cultures of an animal cell strain. *Journal of Biochemical and Microbiological Technology and Engineering*, 11:253–266.
16. Myhr OR and Grong Ø (2000) Modelling of non-isothermal transformations in alloys containing a particle distribution. *Acta Materialia*, Vol. 48(7):1605–1615.
17. Newman K, Buckland ST, Morgan B, King R, Borchers DL, Cole D, Besbeas P, Gimenez O, and Thomas L (2014) *Modelling Population Dynamics: Model Formulation, Fitting and Assessment using State-Space Methods*. Series: Methods in Statistical Ecology, Springer Science+Business Media, New York.
18. Shonkwiler RW and Herod J (2009) *Mathematical Biology: An introduction with Maple and Matlab*. 2nd Edition, Springer Dordrecht, Heidelberg, London, New York.
19. Smoller J (1994) *Shock waves and reaction-diffusion equations*. Springer-Verlag, New York, Second Edition.
20. van Asbrouck J and Taridno P (2009) Using the single seed oxygen consumption measurement as a method of determination of different seed quality parameters for commercial tomato seed samples. *Asian Journal of Food and Agro-Industry*:S88–S95.
21. van Duijn A and Koenig W (2001) Measuring metabolic rate changes. Patent EP1134583 (A1) WO0169243 (A1) US2004033575 (A1) CA2403253 (A1) DE60108480T.
22. Velázquez J, Garrahan JP, and Eichhorn MP (2014) Spatial Complementarity and the Coexistence of Species. *PLoS ONE* 9(12):e114979.
23. Wagner C (1961) Theorie der Alterung von Niederschlägen durch Umlösen (Ostwald-Reifung). *Z. Elektrochem.*, 65:581–591.
24. Xu L, Zhang F, Zhang K, Wang E, and Wang J (2014) The Potential and Flux Landscape Theory of Ecology. *PLoS ONE* 9(1):e86746.